

APPENDIX 5

AMERICAN JOURNAL OF OBSTETRICS AND GYNECOLOGY

ADVISORY COMMITTEE ON POLICY 1953

Willard M. Allen	Nicholson J. Eastman
John L. Brewer	Frederick H. Falls
Francis Bayard Carter	Newell W. Philpott
Conrad G. Collins	John Rock
Samuel A. Cosgrove	Donald G. Tollefson
Walter T. Danneberg	Philip F. Williams

ADVISORY EDITORIAL COMMITTEE 1953

Albert H. Aldridge	Andrew A. Marchetti	Franklin L. Payne
Edward Allen	Harvey B. Matthews	Lawrence M. Randall
Allan C. Barnes	John L. McEldrey	Duncan E. Reid
Leroy A. Calkins	Charles E. McLennan	Ralph A. Reis
Russell R. de Alvarez	Joe Vincent Meigs	Herbert E. Schmitz
R. Gordon Douglas	William F. Meagert	George V. Smith
George H. Gardner	Norman F. Miller	Wm. E. Studdiford
Carl P. Huber	Thaddeus L. Montgomery	E. Stewart Taylor
Frank R. Lock	Daniel G. Morton	Richard W. Te Linde
Curtis J. Lund	Emil Novak	Herbert F. Traut
	Ernest W. Page	

OFFICIAL ORGAN

THE AMERICAN GYNECOLOGICAL SOCIETY; THE AMERICAN ASSOCIATION OF OBSTETRICIANS, GYNECOLOGISTS AND ABDOMINAL SURGEONS; NEW YORK OBSTETRICAL SOCIETY; OBSTETRICAL SOCIETY OF PHILADELPHIA; BROOKLYN GYNECOLOGICAL SOCIETY; ST. LOUIS GYNECOLOGICAL SOCIETY; NEW ORLEANS GYNECOLOGICAL AND OBSTETRICAL SOCIETY; THE OBSTETRICAL AND GYNECOLOGICAL SOCIETY OF MARYLAND; CHICAGO GYNECOLOGICAL SOCIETY; CINCINNATI OBSTETRICAL SOCIETY; CENTRAL ASSOCIATION OF OBSTETRICIANS AND GYNECOLOGISTS; AMERICAN BOARD OF OBSTETRICS AND GYNECOLOGY; WASHINGTON GYNECOLOGICAL SOCIETY; PITTSBURGH OBSTETRICAL AND GYNECOLOGICAL SOCIETY; OBSTETRICAL SOCIETY OF BOSTON; LOUISVILLE OBSTETRICAL AND GYNECOLOGICAL SOCIETY; SOUTH ATLANTIC ASSOCIATION OF OBSTETRICIANS AND GYNECOLOGISTS; SEATTLE GYNECOLOGICAL SOCIETY; SOCIETY OF OBSTETRICIANS AND GYNECOLOGISTS OF CANADA; ALABAMA ASSOCIATION OF OBSTETRICIANS AND GYNECOLOGISTS; ANKON OBSTETRICAL AND GYNECOLOGICAL SOCIETY; KANSAS CITY GYNECOLOGICAL SOCIETY; CENTRAL NEW YORK ASSOCIATION OF GYNECOLOGISTS AND OBSTETRICIANS; NEW JERSEY OBSTETRICAL AND GYNECOLOGICAL SOCIETY; IOWA OBSTETRICAL AND GYNECOLOGICAL SOCIETY; THE TEXAS ASSOCIATION OF OBSTETRICIANS AND GYNECOLOGISTS; OKLAHOMA CITY OBSTETRICAL AND GYNECOLOGICAL SOCIETY; MEMPHIS OBSTETRICAL AND GYNECOLOGICAL SOCIETY; UTAH OBSTETRICAL AND GYNECOLOGICAL SOCIETY; ROCHESTER OBSTETRICAL AND GYNECOLOGICAL SOCIETY

EDITORS

HOWARD C. TAYLOR, JR., and WILLIAM J. DIECKMANN

ADVISORY EDITOR
GEORGE W. KOSMAK

VOLUME 66
JULY—DECEMBER, 1953

THE C. V. MOSBY COMPANY
ST. LOUIS
1953

Volume 66
Number 5DOES THE ADMINISTRATION OF DIETHYLSTILBESTROL DURING PREGNANCY HAVE THERAPEUTIC VALUE?^{1,2}W. J. DIECKMANN, M.D., M. E. DAVIS, M.D., L. M. RYNKIEWICZ, S.M., AND
R. E. PORTINGER, S.M., CHICAGO, ILL.*(From the Department of Obstetrics and Gynecology of the University of Chicago and the Chicago Lying-in Hospital)*

IN 1946 Smith and Smith¹ suggested that increasing amounts of diethylstilbestrol should be administered to all women during pregnancy to prevent or decrease the hazards of the late complications of pregnancy for mothers and babies. The basis for such prophylactic therapy as well as the active therapy of these pregnancy complications stems from a series of experiments by the Smiths on the steroid hormones in normal and abnormal pregnancy. These laboratory observations and their theoretical implications were supported by clinical observations, part of which were made under the supervision of the Smiths and part were the collected reports of other clinical observers.

The use of diethylstilbestrol to prevent and to treat pregnancy complications is based on the supposition that there develops a deficiency in the production of progesterone and other steroids by the placenta which predisposes to or causes these pregnancy complications. The secretion of these steroids can be stimulated by diethylstilbestrol. The increased amounts of steroids made available by the placenta postpone, reduce the severity of, or prevent some of the late complications of pregnancy.

The laboratory experiments which provided the background for this interesting concept of the Smiths have lacked confirmation by other investigators. Davis and Fugo² in two reports noted that the administration of diethylstilbestrol to patients during pregnancy did not result in an increased output of urinary pregnanediol, a measure of progesterone metabolism. Sommerville, Marrian and Clayton³ confirmed these observations and noted a drop in urinary pregnanediol and no gross change in endogenous estrogen. Although many additional experimental data will be necessary to determine the role of diethylstilbestrol in placental steroid metabolism, this paper will confine itself to the clinical implications of the Smith concept.

Smith and Smith in 1949⁴ reported on the influence of diethylstilbestrol on the progress and outcome of pregnancy in a series of primigravidas. As

¹This investigation was supported in part by a research grant PHS RG3570 from the National Institutes of Health, Public Health Service.

²Presented at the Seventy-sixth Annual Meeting of the American Gynecological Society, Lake Placid, N. Y., June 15 to 17, 1953.

controls they used a
They recorded the f
the late toxemias of
usually large for th
was decreased. (4)
decreased. (5) The r

The most serious
Patients to whom th
meticulous study and
ously. In two nutrit
tients who were coor
instructions, keep go
dence of abortion, pr
a lower perinatal dea
women delivered con
were compared with
received identical tre
significant differences

The prophylactic
use in pregnancy com
to postpone, or to an
mothers and babies i
plished by the daily co
a careful perusal of
not supported by adeq

The properly con
should have no knowl
of patients should re
the medication on tri
treated simultaneously
cently reported by F
that diethylstilbestrol
maturity, perinatal m

We felt that it w
clinical experiment to
practice. The first pa
study terminated Nov.

The following cri
large so that the resu
of patients who were o
was decided that 2,000
prior to the twentieth
Every other patient w
sonal element, each pr
individual who was no

Since our data were at variance with those of the Smiths, they were all rechecked. The charts of patients with toxemia of pregnancy, premature delivery, stillbirths and neonatal deaths, and any other complication or abnormality, were examined again by one of the senior authors with no knowledge of the kind of medication. There was no significant change in any of the results.

TABLE IX. CONGENITAL ANOMALIES

TYPE OF ANOMALY	PRIMIPARAS		MULTIPARAS	
	STILBESTROL	CONTROL	STILBESTROL	CONTROL
Minor	7	7	9	4
Skin, as papilloma	7	12	7	6
Cystocele, hydrocele	4	3	3	2
Harelip, cleft palate, etc.	1	0	0	1
Clubfeet, multiple digits	2	5	6	2
Mongolism	0	0	0	1
Brain and spinal cord	1	0	0	0
Cardiac, etc.	2	1	1	2
Gastrointestinal	1	0	0	0
Genitourinary	0	2	0	3
Multiple major	2	2	1	1
Total anomalies	27	32	27	24
Total infants	426	415	375	351

Conclusions

A strictly controlled clinical trial of the therapeutic value of diethylstilbestrol administered to patients during pregnancy in reducing the hazards of some of the late complications of pregnancy for mothers and babies has been reported.

The various complications were studied in the total unselected group of patients divided into primigravidas, primiparas, and multiparas. Then the groups were again studied after all groups were corrected to compare with the Smiths'.

The results of the administration of diethylstilbestrol in graduated amounts to 840 patients according to a schedule suggested by the Smiths were compared with the results of an identical placebo tablet given to 806 patients. Stilbestrol did not reduce the incidence of abortion, prematurity, or postmaturity. Premature babies of stilbestrol-treated mothers were no longer nor more mature for their gestational ages than comparable prematures in the control group of placebo-treated mothers. It did not decrease the incidence of perinatal mortality. It did not decrease the frequency of the toxemias of pregnancy.

Acknowledgment is made to Eli Lilly and Company for aid in making the stilbestrol and placebo tablets with the dye and for the final determination of the stilbestrol; to Lillian Natusko for the examination of the urines for phenol red; to the staff and residents for their cooperation.

References

1. Smith, O. W., and Smith, G. van S.: *Am. J. Obst. & Gynec.* 51: 411, 1946.
2. Smith, G. van S., and Smith, O. W.: *Physiol. Rev.* 28: 1, 1948.
3. Davis, M. Edward, and Fugo, N. W.: *Proc. Soc. Exper. Biol. & Med.* 65: 283, and 56: 391, 1947.

4. Davis, M. E.
5. Sommerville,
6. Smith, O. W.
7. Dieckmann,
8. Gitman, L.,
9. Smith, O. W.
10. Ferguson, J.
11. Cacario, E. A.

DR. GEORGE
mann and his ass
work, and exper
told us he was ge
for a beneficial e
value could be r
planning to anal
Worcester, Assist
who had had cor
nuancy, stated th
thousands of each
She advised that
every reason to
treated group wh
published by Dr.
studied. To bre
impossible. Ther
We too are disap
to so much troubl
run, their experim
We also wis
that the only diff
groups lay in the
teacher, who saw
dentally, fully an
drawal in patien
conveyed this con
Dr. Olive Smith, t
Actually, most of
and, therefore, ma

Neither the
patients was set c
omitted from our
those who had thr
our control series.
who began taking
phyliactic value of
previous abortions
peatedly miscarry
now are certain th
ing abortion, there
They include
There were more
trols (23 vs. 17) a
pre-eclampsia. If